

## REFERENCES

Anderson, E.H., Ward, L.R., De Sax, M.J., and DeSa, J.D.H. (1977). Bacteriophage – Typing Designations of *S. typhimurium*. *Journal of Hygiene* **78**, 297 – 300.

Defra (2005) Zoonoses Report UK 2004 (in press).

EEC (1997) European Union Council Directive 92/117/EEC and amendment 97/22/EC. *Official Journal* L062, pp 38 – 48.

EEC (2003) Zoonoses Regulations 2160/2003.

Evans, S.J., Davies, R.H., Binns, S.H., Liebana, E., Jones, T.W.H., Millar, M.F., Threlfall, E.J., Ward, L.R., Hopkins, K.L., Mackay, P.H.S., and Gayford, P.J.R. (2005), Multiple antimicrobial resistant *Salmonella enterica* serovar Paratyphi B variant Java in cattle: a case report. *Veterinary Record* **156**, (11), 343 – 346.

HMSO (1989) The Zoonoses Order 1989 (ISBN 0 11 096285 0).

HMSO (1993) The Poultry Breeding Flocks and Hatcheries Order 1993 (ISBN 0 11 034898 2).

HMSO (2002) The TSE Regulations 2002 (ISBN 0 11 039914 5).

HMSO (2003) The Animal By-Products Regulations 2003 (ISBN 0 11 046674 8).

Le Minor, L., Popoff, M.Y. (1987) Designation of *Salmonella enterica* sp. nov., nom. rev., as the type and only species of the genus *Salmonella*. *International Journal of Systematic Bacteriology* **37**, 465-468.

VLA (2005) Veterinary Investigation Surveillance Report (VIDA) 2004, (ISBN 1 8995 1325 6).

Ward, L.R., DeSa, J.D.H. and Rowe, B. (1987). A Phage typing Scheme for *Salmonella enteritidis*. *Epidemiology and Infection*, **99**, 291 – 294.

## FURTHER PUBLICATIONS OF INTEREST

- Carroll P; La Ragione RM Sayers AR; Woodward MJ (2004). The O-antigen of *Salmonella enterica* serotype Enteritidis PT4: a significant factor in gastrointestinal colonisation of young but not newly hatched chicks. *Veterinary Microbiology* **102** (1-2) 73-85.
- Cogan TA; Jorgensen F; Lappin-Scott HM; Benson CE; Woodward MJ; Humphrey TJ (2004). Flagella and curli fimbriae are important for the growth of *Salmonella enterica* serovars in hen eggs. *Microbiology* **150** (4) 1063-1071.
- Coldham NG Woodward MJ (2004). Characterization of the *Salmonella* Typhimurium proteome by semi-automated two dimensional HPLC-mass spectrometry: detection of proteins implicated in multiple antibiotic resistance." *Journal of Proteome Research* **3** (3) 595-603.
- Cook AJC (2004). Measuring the impact of *Salmonella* control in finishing pigs: lessons from a pilot study. *Pig Journal* (53) 157-163.
- Davies, R.H., Paiba, G., Evans, S.J., Dalziel, R. (2000) Surveys for *Salmonella* in pigs, cattle and sheep at slaughter in Great Britain. *Veterinary Record*, **147**, 695.
- Davies R.H., Dalziel R., Wilesmith J.W., Ryan J., Evans S.J., Paiba G.A., Byrne C., Pascoe S. (2001) National survey for *Salmonella* in pigs at slaughter in Great Britain. Proceedings of the 4<sup>th</sup> International symposium on the Epidemiology and Control of *Salmonella* and other Foodborne Pathogens in Pork. 2-5 September 2001, Leipzig, Germany, 162-173
- Davies RH; Dalziel R; Gibbens JC; Wilesmith JW; Ryan JMB; Evans SJ; Byrne C; Paiba GA; Pascoe SJS; Teale CJ (2004). National survey for *Salmonella* in pigs, cattle and sheep at slaughter in Great Britain (1999-2000). *Journal of Applied Microbiology* **96** (4) 750-760.
- Davies R; Breslin M (2004). Observations on salmonella contamination of eggs from infected commercial laying flocks where vaccination for *Salmonella enterica* serovar Enteritidis had been used. *Avian Pathology* **33** (2) 133-144.
- Davies R (2004). Fowl typhoid and pullorum disease. In: Manual of diagnostic tests and vaccines for terrestrial animals (mammals, birds and bees), 5th ed, OIE, 2004, 868-877.

Defra (1989a) Code of practice for the control of *Salmonella* during the storage, handling and transport of raw materials intended for incorporation into, or direct use as, animal feedingstuffs.

Defra (1989b) Code of practice for the control of *Salmonella* in the production of final feed for livestock in premises producing less than 10,000 tonnes per annum.

Defra (1989c) Code of practice for the control of *Salmonella* in the production of final feed for livestock in premises producing over 10,000 tonnes per annum.

Defra (1989d) Code of practice for the control of *Salmonella* in the animal by-products rendering industry.

Defra (1989e) Code of practice for the control of *Salmonella* for the UK fish meal industry.

Defra (2001) Code of practice for the prevention and control of *Salmonella* on pig farms.

Evans, S. (2000) *Salmonella* monitoring and control in cattle and sheep. In: Zoonotic infections in livestock and the risk to public health. Proceedings of a meeting organised by MAFF and held on 7<sup>th</sup> December 2000, London, UK.

Gradel KO; Sayers AR; Davies RH (2004). Surface disinfection tests with *Salmonella* and a putative indicator bacterium, mimicking worst-case scenarios in poultry houses. *Poultry Science* **83** (10) 1636-1643.

Humphreys S; Rowley G; Stevenson A; Anjum MF; Woodward MJ; Gilbert S; Kormanec J; Roberts M (2004). Role of the two-component regulator CpxAR in the virulence of *Salmonella enterica* serotype Typhimurium. *Infection and Immunity* **72** (8) 4654-4661.

Ivanek R; Snary EL; Cook AJC; Grohn YT (2004). A mathematical model for the transmission of *Salmonella* Typhimurium within a grower-finisher pig herd in Great Britain. *Journal of Food Protection* **67** (11) 2403-2409.

La Ragione RM; Narbad A; Gasson MJ; Woodward MJ (2004). In vivo characterization of *Lactobacillus johnsonii* FI9785 for use as a defined competitive exclusion agent against bacterial pathogens in poultry. *Letters in Applied Microbiology* **38** (3) 197-205.

Liebana E; Clouting C; Garcia-Migura L; Clifton-Hadley Fa; Lindsay E; Threlfall EJ; Davies RH (2004). Multiple genetic typing of *Salmonella*

Enteritidis phage types 4, 6, 7, 8 and 13a isolates from animals and humans in the UK. *Veterinary Microbiology* **100** (3-4) 189-195.

Liebana E; Batchelor M; Torres C; Brinas L; Lagos LA; Abdalhamid B; Hanson ND; Martinez-Urtaza J (2004). Pediatric infection due to multiresistant *Salmonella enterica* serotype Infantis in Honduras. *Journal of Clinical Microbiology* **42** (10) 4885-4888.

Liebana E; Gibbs M; Clouting C; Barker L; Clifton-Hadley FA; Pleydell E; Abdalhamid B; Hanson ND; Martin L; Poppe C; Davies RH (2004). Characterization of beta-lactamases responsible for resistance to extended-spectrum cephalosporins in *Escherichia coli* and *Salmonella enterica* strains from food-producing animals in the United Kingdom. *Microbial Drug Resistance* **10** (1) 1-9.

Martinez-Urtaza J; Liebana E; Garcia-Migura L; Perez-Pineiro P; Saco M (2004). Characterization of *Salmonella enterica* serovar Typhimurium from marine environments in coastal waters of Galicia (Spain). *Applied and Environmental Microbiology* **70** (7) 4030-4034.

Randall LP; Cooles SW; Osborn MK; Piddock LJV; Woodward MJ (2004). Antibiotic resistance genes, integrons and multiple antibiotic resistance in thirty-five serotypes of *Salmonella enterica* isolated from humans and animals in the UK. *Journal of Antimicrobial Chemotherapy* **53** (2) 208-216.

Randall LP; Cooles SW; Piddock LJV; Woodward MJ (2004). Effect of triclosan or a phenolic farm disinfectant on the selection of antibiotic-resistant *Salmonella enterica*. *Journal of Antimicrobial Chemotherapy* **54** (3) 621-627.

Thomson N; Baker S; Pickard D; Fookes M; Anjum M; Hamlin N; Wain J; House D; Bhutta Z; Chan K; Falkow S; Parkhill J; Woodward M; Ivens A; Dougan G (2004). The role of prophage-like elements in the diversity of *Salmonella enterica* serovars." *Journal of Molecular Biology* (339) 279-300.

Wray C; Davies R (2004). *Salmonella* in cattle: update. *Cattle Practice* **12** (4) 313-316.

## Acknowledgements

Incidents were reported by Nominated Officers of the Veterinary Laboratories Agency for England and Wales and Divisional Veterinary Managers for Scotland and, through them, by private laboratories.

Divisional Veterinary Managers of the State Veterinary Service are responsible for the collection of samples of processed animal protein.

Staff of the Veterinary Laboratories Agency processed the data.

The following reference laboratories made or confirmed the majority of isolations:

- Veterinary Laboratories Agency, Weybridge and Lasswade.
- HPA Laboratory of Enteric Pathogens, Colindale.
- Scottish *Salmonella* Reference Laboratory, Glasgow.

This report was compiled by:

Sue Kidd and Christina Papadopoulou  
Centre for Epidemiology and Risk Analysis,  
Veterinary Laboratories Agency,  
New Haw,  
Addlestone,  
Surrey,  
KT15 3NB

Telephone: + 44 (0)1932 - 357622

Fax: + 44 (0)1932 - 349983

E-mail [s.a.kidd@vla.defra.gsi.gov.uk](mailto:s.a.kidd@vla.defra.gsi.gov.uk)

Further copies of the report are available from the above address

VLA is an Executive Agency of the Department for Environment, Food and Rural Affairs.